


# New Heavy Quark Baryons

Michal Kreps

University of Karlsruhe



# Why heavy baryon spectroscopy

- Heavy Quark mesons are QCD analog of "hydrogen atom"
  - Starts to be very sensitive test of various model in non-perturbative regime of QCD
  - Lot of information in charm sector
  - Bottom sector starts to speak up as well
- Heavy Quark baryon are next interesting laboratory
  - Heavy quark - light diquark is basic picture
  - Another sensitive test of models
  - Still many things to observe in charm sector
  - In bottom sector only  $\Lambda_b$  directly seen
- Discovery of new particles is exciting and fun

# Where to study heavy baryons



- Everywhere where we produce them and have detector to detect them
- Current results come from
  - B-factories (Belle, *BABAR*)
    - + Have large amount of data
    - + Clean environment
    - Bound to charm sector
  - Tevatron (CDF)
    - Difficult environment from  $p\bar{p}$  collisions
    - Only now starts to have reasonable amount of data for b-baryons
    - + Can do all b-hadrons

# Directly observed states

Listed in PDG 2006

Not in PDG 2006, covered here

Listed in PDG 2006, but new results







# $\Lambda_c^+(2880), \Lambda_c^+(2940)$



- $287 \text{ fb}^{-1}$  of data
- $p D^0$  final state
- $D^0 \rightarrow K\pi, D^0 \rightarrow K\pi\pi\pi$
- PRL 98, 012001 (2007)
- $553 \text{ fb}^{-1}$  of data
- Confirmation in  $\Lambda_c^+ \pi^+ \pi^-$
- $\Lambda_c^+ \pi^\pm$  consistent with  $\Sigma_c(2455)$
- hep-ex/0608043





# $\Lambda_c^+(2880), \Lambda_c^+(2940)$


- $\Lambda_c(2880)$  known state, but  $pD^0$  decay is new
- $\Lambda_c(2940)$  observed for the first time
- Significance  $7.5\sigma$  at *BABAR* and  $6.2\sigma$  at Belle
- Mass and width consistent between experiments

	State	Mass [MeV/c <sup>2</sup> ]	Width [MeV/c <sup>2</sup> ]
<i>BABAR</i>	$\Lambda_c(2880)$	$2882 \pm 0.1 \pm 0.5$	$5.8 \pm 1.5 \pm 1.1$
Belle	$\Lambda_c(2880)$	$2881.2 \pm 0.2 \pm 0.4$	$5.5 \pm 0.7 \pm 1.1$
<i>BABAR</i>	$\Lambda_c(2940)$	$2939.8 \pm 1.3 \pm 1.0$	$17.5 \pm 5.2 \pm 5.9$
Belle	$\Lambda_c(2940)$	$2938.0 \pm 1.3^{+2.0}_{-4.0}$	$13^{+8}_{-5}{}^{+27}_{-7}$

- To learn more, both experiments do further studies
  - *BABAR* checks isospin partners
  - Belle studies resonant substructure of decay and angular distributions



# $\Lambda_c^+(2880), \Lambda_c^+(2940)$




Curves same rate as  $pD^0$

- If  $\Sigma_c \Rightarrow$  also  $\Sigma_c^{++} \rightarrow D^+ p$   
 $D^+ \rightarrow K\pi\pi$
- No resonant structure seen
- ⇒ Both states are  $\Lambda_c$ 's
  - 3  $\Lambda_c$  states predicted  $\approx 2940$  MeV/ $c^2$   
 $J^P = (1/2)^+, (1/2)^-, (3/2)^-$   
Migura et al, Eur.Phys.J. A28 (2006) 41
  - The  $\Lambda_c(2880)^+$  is near a predicted  $(3/2)^-$  state.
  - Details PRL 98, 012001 (2007)

# $\Lambda_c^+(2880)$



Fit  $\Lambda_c(2880)$  yield in bins of  $M(\Lambda_c^+\pi^\pm)$

Details: hep-ex/0608043



- Significance of  $\Lambda_c(2880) \rightarrow \Sigma_c(2520)\pi$   $3\sigma$  with syst.
- $\Gamma(\Sigma_c(2455)\pi)/\Gamma(\Lambda_c\pi\pi) = 40.4 \pm 2.1 \pm 1.4\%$
- $\Gamma(\Sigma_c(2520)\pi)/\Gamma(\Lambda_c\pi\pi) = 9.1 \pm 2.5 \pm 1.0\%$
- $\Gamma(\Sigma_c(2520)\pi)/\Gamma(\Sigma_c(2455)\pi) = 22.5 \pm 6.2 \pm 2.5\%$

# $\Lambda_c^+(2880)$



- Fit  $\Lambda_c(2880)$  mass distribution in angular bins and subtract non-resonant contribution
- $\chi^2/ndf.$ : 46.7/9 ( $J = 1/2$ ); 35.1/8 ( $J = 3/2$ ); 12.1/7 ( $J = 5/2$ )
- From  $\chi^2$  difference exclude  $J = 1/2$  ( $J = 3/2$ ) by  $5.5\sigma$  ( $4.8\sigma$ )
- HQS expectations for  $\Gamma(\Sigma_c(2520)\pi)/\Gamma(\Sigma_c(2455)\pi)$ :  
140% ( $J^P = 5/2^-$ ) and 23 – 36% ( $J^P = 5/2^+$ )

# $\Xi_c(2980), \Xi_c(3077)$

- Belle searches for new states in  $\Lambda_c^+ K^- \pi^+$  and  $\Lambda_c^+ K_s^0 \pi^+$
- Uses  $462 \text{ fb}^{-1}$  of data
- $\Lambda_c^+ \rightarrow p K^- \pi^+$  decay
- Signal events:

$\Xi_c^+(2980)$   $405.3 \pm 50.7$

$\Xi_c^+(3077)$   $326.0 \pm 39.6$

$\Xi_c^0(2980)$   $42.3 \pm 23.8$

$\Xi_c^0(3077)$   $67.1 \pm 19.9$



- Significance:

$\Xi_c^+(2980)$   $5.7\sigma$

$\Xi_c^+(3077)$   $9.2\sigma$

$\Xi_c^0(2980)$   $1.5\sigma$

$\Xi_c^0(3077)$   $4.4\sigma$





Details in PRL 97, 162001 (2006)



# $\Xi_c(2980), \Xi_c(3077)$


- Using  $289 \text{ fb}^{-1}$  of data *BABAR* confirms Belle's observation
- Study only  $\Lambda_c^+ K^- \pi^+$  with  $\Lambda_c^+ \rightarrow p K^- \pi^+$  decays
- Observes two structures in mass difference spectra
- Number of signal events
  - $\Xi_c^+(2980) \quad 284 \pm 45 \pm 46$
  - $\Xi_c^+(3077) \quad 204 \pm 35 \pm 12$
- Significance
  - $\Xi_c^+(2980) \quad 7.0 \sigma$
  - $\Xi_c^+(3077) \quad 8.6 \sigma$
- Details in [hep-ex/0607042](https://arxiv.org/abs/hep-ex/0607042)







# $\Xi_c(2980), \Xi_c(3077)$


$2.95 < M_{\Xi_c} < 2.99 \text{ GeV}$



$3.07 < M_{\Xi_c} < 3.09 \text{ GeV}$






$3.02 < M_{\Xi_c} < 3.05 \text{ GeV}$



$3.12 < M_{\Xi_c} < 3.22 \text{ GeV}$



# $\Xi_c(2980), \Xi_c(3077)$



- 2D fit to  $M(\Lambda_c^+\pi^+)$  and  $M(\Lambda_c^+K^-\pi^+)$
- Fit for resonance substructure of  $\Xi_c$  decays
- Allow for  $\Sigma_c$  states also in background



# $\Xi_c(2980), \Xi_c(3077)$

- *BABAR* and Belle measure consistent masses and widths

	State	Mass [MeV/c <sup>2</sup> ]	Width [MeV/c <sup>2</sup> ]
Belle	$\Xi_c^0(2980)$	$2977.1 \pm 8.8 \pm 3.5$	43.5 (fixed)
Belle	$\Xi_c^+(2980)$	$2978.5 \pm 2.1 \pm 2.0$	$43.5 \pm 7.5 \pm 7.0$
<i>BABAR</i>	$\Xi_c^+(2980)$	$2967.1 \pm 1.9 \pm 1.0$	$23.6 \pm 2.8 \pm 1.3$
Belle	$\Xi_c^0(3077)$	$3082.8 \pm 1.8 \pm 1.5$	$5.2 \pm 3.1 \pm 1.8$
Belle	$\Xi_c^+(3077)$	$3076.7 \pm 0.9 \pm 0.5$	$6.2 \pm 1.2 \pm 0.8$
<i>BABAR</i>	$\Xi_c^+(3077)$	$3076.4 \pm 0.7 \pm 0.3$	$6.2 \pm 1.6 \pm 0.5$



- *BABAR* adds resonant substructure





$\Xi_c^+(2980) \rightarrow \Sigma_c^{++}(2455)K^-$	$132 \pm 31 \pm 5$	$4.9\sigma$
$\Xi_c^+(2980) \rightarrow \Lambda_c^+K^-\pi^+$	$152 \pm 37 \pm 45$	$4.1\sigma$
$\Xi_c^+(3077) \rightarrow \Sigma_c^{++}(2455)K^-$	$87 \pm 20 \pm 4$	$5.8\sigma$
$\Xi_c^+(3077) \rightarrow \Sigma_c^{++}(2520)K^-$	$82 \pm 23 \pm 6$	$4.6\sigma$
$\Xi_c^+(3077) \rightarrow \Lambda_c^+K^-\pi^+$	$35 \pm 24 \pm 1$	$1.4\sigma$




$$\Xi_c' \rightarrow \Xi_c \gamma$$

- State seen by CLEO in 1999 [PRL 82, 492(1999)]
- Seen and studied in  $\Xi_c' \rightarrow \Xi_c \gamma$  decay
- Not confirmed since then
- It is lightest state above ground state
- BABAR uses both  $c\bar{c}$  events and  $B$  decays of  $232 \text{ fb}^{-1}$  of data
- $\Xi_c^+ \rightarrow \Xi^- \pi^+ \pi^+$
- $\Xi_c^0 \rightarrow \Xi^- \pi^+$
- Details in hep-ex/0607086







- $c\bar{c}$  cross section

$$\sigma(e^+e^- \rightarrow \Xi_c'^+ X) \times \mathcal{B}(\Xi_c'^+ \rightarrow \Xi^- \pi^+ \pi^+) = 141 \pm 24 \pm 19 \text{ fb}$$

$$\sigma(e^+e^- \rightarrow \Xi_c'^0 X) \times \mathcal{B}(\Xi_c'^0 \rightarrow \Xi^- \pi^+) = 70 \pm 11 \pm 6 \text{ fb}$$

$$\sigma(e^+e^- \rightarrow \Xi_c^0 X) \times \mathcal{B}(\Xi_c^0 \rightarrow \Xi^- \pi^+) = 388 \pm 39 \pm 41 \text{ fb}$$



- $B$  production rate

$$\mathcal{B}(B \rightarrow \Xi_c'^+ X) \times \mathcal{B}(\Xi_c'^+ \rightarrow \Xi^- \pi^+ \pi^+) = (1.69 \pm 0.17 \pm 0.10) \cdot 10^{-4}$$

$$\mathcal{B}(B \rightarrow \Xi_c'^0 X) \times \mathcal{B}(\Xi_c'^0 \rightarrow \Xi^- \pi^+) = (0.67 \pm 0.07 \pm 0.03) \cdot 10^{-4}$$

$$\mathcal{B}(B \rightarrow \Xi_c^0 X) \times \mathcal{B}(\Xi_c^0 \rightarrow \Xi^- \pi^+) = (2.11 \pm 0.019 \pm 0.25) \cdot 10^{-4}$$

- From helicity angle  $J = 1/2$ , higher spin cannot be excluded






$$\Omega_c^* \rightarrow \Omega_c \gamma$$

- $\Omega_c^*$  last unobserved ground state charm baryon
- *BABAR* search uses  $231 \text{ fb}^{-1}$  data
- $\Omega_c$  reconstructed in channels:  
 $\Omega^- \pi^+; N = 156 \pm 15$  (a)  
 $\Omega^- \pi^+ \pi^0; N = 92 \pm 26$  (b)  
 $\Omega^- \pi^+ \pi^- \pi^+; N = 23 \pm 10$  (c)  
 $\Xi^- K^- \pi^+ \pi^+; N = 34 \pm 15$  (d)  
 $\Omega^- \rightarrow \Lambda K^-, \Xi^- \rightarrow \Lambda \pi^-, \Lambda \rightarrow p \pi^-$
- Combine  $\Omega_c$  with  $\gamma$


Details in PRL 97, 232001 (2006)





$$\Omega_c^* \rightarrow \Omega_c \gamma$$

- Use  $\Delta M = M_{\Omega_c \gamma} - M_{\Omega_c}$  for better resolution
  - (a)  $\Delta M = 69.9 \pm 1.4 \pm 1.0 \text{ MeV}/c^2$   
 $N = 39^{+10}_{-9} \pm 6$ ; significance  $4.2\sigma$
  - (b)  $\Delta M = 71.8 \pm 1.3 \pm 1.1 \text{ MeV}/c^2$   
 $N = 55^{+16}_{-15} \pm 6$ ; significance  $3.4\sigma$
  - (c)  $\Delta M = 69.9 \text{ MeV}/c^2$  (fixed)  
 $N = -5 \pm 5 \pm 1$ ;
  - (d)  $\Delta M = 69.4^{+1.9}_{-2.0} \pm 1.0 \text{ MeV}/c^2$   
 $N = 20 \pm 9 \pm 3$ ; significance  $2.0\sigma$
- Channels consistent  
 ⇒ combine them






$$\Omega_c^* \rightarrow \Omega_c \gamma$$

- Fit with Crystal ball function for signal with fixed shape parameters
- $\Delta M = 70.8 \pm 1.0 \pm 1.1 \text{ MeV}/c^2$
- $N = 105 \pm 21 \pm 6$
- $5.2\sigma$  significance

Predictions for  $\Delta M$  ( $\text{MeV}/c^2$ )

- HQET 80
- Lattice QCD 90
- Non-relativistic Quark Model 50-73



# $\Sigma_b$ expectations

- Up to now  $\Lambda_b$  only directly observed  $b$ -baryon
- Lack of the experimental result mainly due to the statistics
- Tevatron experiments start to have enough statistics to search for other  $b$ -baryons
- Decay via p-wave  $\pi$

$$\Sigma_b: bqq \quad J^P = S_Q + S_{qq}$$

$$\rightarrow 3/2^+ (\Sigma_b^*)$$

$$\rightarrow 1/2^+ (\Sigma_b)$$


$\Sigma_b^{(*)+}$	$uub$
$\Sigma_b^{*-}$	$ddb$
$\Sigma_b^{*0}$	$udb$

$\Sigma_b^{*0}$  decays through  $\pi^0$ ,  
can't be seen at CDF

Property	Expectation (MeV/c <sup>2</sup> )
$m(\Sigma_b) - m(\Lambda_b)$	180 — 210
$m(\Sigma_b^*) - m(\Sigma_b)$	10 — 40
$m(\Sigma_b^-) - m(\Sigma_b^+)$	5 — 7
$\Gamma(\Sigma_b), \Gamma(\Sigma_b^*)$	$\approx 8, \approx 15$


# $\Lambda_b$ and sample composition

CDF II Preliminary,  $L = 1.1 \text{ fb}^{-1}$




$\Lambda_b$	86%
$B$ mesons	10%
combinatorial	4%

- $\Lambda_b \rightarrow \Lambda_c^+ \pi^-, \Lambda_c^+ \rightarrow p K^- \pi^+$
- With  $1.1 \text{ fb}^{-1}$  around 3000  $\Lambda_b \Rightarrow$  world's largest sample
- For  $\Sigma_b$  search select narrow region around fully reconstructed peak



# Background estimation

- Do blind search
- Fix all backgrounds before looking to signal region
- Shapes  
 $\Lambda_b$  sideband  
PYTHIA MC  
B miss-reconstructed data
- Relative normalization according to  $\Lambda_b$  mass fit
- Determined background describes data well




# Unblinded $\Sigma_b$ Q-distribution



- Excess after unblinding

	Data	bkg	excess
$\Sigma_b^{(*)-}$	416	268	148
$\Sigma_b^{(*)+}$	406	298	108

- Data indicate two peaks for each charge
- Do unbinned maximum likelihood fit
- Fit for Q values and number of events



# $\Sigma_b$ Fit result

- Mass differences ( $\text{MeV}/c^2$ )

$$\rightarrow m(\Sigma_b^-) - m(\Lambda_b) - m(\pi) = 55.9 \pm 1.0(\text{stat}) \pm 0.1(\text{sys})$$

$$\rightarrow m(\Sigma_b^+) - m(\Lambda_b) - m(\pi) = 48.4^{+2.0}_{-2.3}(\text{stat}) \pm 0.1(\text{sys})$$

$$\rightarrow m(\Sigma_b^*) - m(\Sigma_b) = 21.3^{+2.0}_{-1.9}(\text{stat})^{+0.4}_{-0.2}(\text{sys})$$


- Signal events

$$\rightarrow N(\Sigma_b^+) = 29^{+12.4}_{-11.6}(\text{stat})^{+5.0}_{-3.4}(\text{sys})$$

$$\rightarrow N(\Sigma_b^-) = 60^{+14.8}_{-13.8}(\text{stat})^{+8.4}_{-4.0}(\text{sys})$$

$$\rightarrow N(\Sigma_b^{*+}) = 74^{+17.2}_{-16.3}(\text{stat})^{+10.3}_{-5.7}(\text{sys})$$

$$\rightarrow N(\Sigma_b^{*-}) = 74^{+18.2}_{-17.4}(\text{stat})^{+15.6}_{-5.0}(\text{sys})$$



# $\Sigma_b$ Significance

- Repeat fit with alternative hypothesis
  - Single peak left out
  - Only one peak in each charge combination
  - No peak, pure background
- Derived from  $\Delta(-\ln \mathcal{L})$

Hypothesis	$\Delta(-\ln \mathcal{L})$	Hypothesis	$\Delta(-\ln \mathcal{L})$
Null	44.7	No $\Sigma_b^-$	10.4
2 peaks	14.3	No $\Sigma_b^+$	1.1
		No $\Sigma_b^{*-}$	10.1
		No $\Sigma_b^{*+}$	9.8

- ⇒ Significance more than  $5\sigma$  for 4 peaks
- ⇒ Evidence for three out of four individual peaks

Details at

<http://www-cdf.fnal.gov/physics/new/bottom/060921.blessed-sigmab>

# Conclusions

- Last year very rich for Heavy Quark Baryons
  - Several new baryon states in charm sector discovered  
 $\Lambda_c^+(2940)$ ,  $\Xi_c^{+,0}(2980)$ ,  $\Xi_c^{+,0}(3077)$  and  $\Omega_c^*$
  - Several refined measurements in charm sector
  - Charged  $\Sigma_b$  states discovered in bottom sector
- ⇒ Our knowledge about Heavy Quark Baryons increased
- I'm convinced this was not our last word on the topic